


TRIGGER FINGER


The **flexor tendons** that bend the fingers and thumb run from the forearm. The tendons run within a tube called the **flexor sheath**. This has a complex system of thickenings called “annular” pulleys. There are five in the fingers (A1-A5) and three in the thumb (A1, oblique and A2). The whole system is mechanically very similar to bicycle brake cable.



The tendons are lined by a layer of tissue called the **tenosynovium**. This allows them to glide smoothly through the various pulleys. As we age, the tenosynovium degenerates and thickens, a condition called **tenosynovitis**. This in alone can make the fingers feel puffy, tender, “slow” and even stiff.


Trigger finger is caused by the thickening around and on the tendon, which causes it to catch as it runs in and out of the sheath. You can often feel this swelling in the palm as you move the finger. The condition can occur in any and sometimes multiple fingers. There are two ways of treating the problem.

Steroid injection A small amount of steroid is injected around the tendon. This flattens out the swelling on the tendon allowing it to glide freely in and out of the sheath once more. A single injection is all that is needed in 50% of cases. A further 25% will respond to another injection (i.e. three-quarters of cases can be successfully treated in this way). The steroid injection does not work immediately. It causes no general side-effects but occasionally the skin around the injection-site can be made a little thinner. Therefore two injections are the maximum for the treatment of each “episode”. Even after successful treatment, the triggering may return in the affected or other fingers. This is, however, very unusual if you have had surgery



Surgery This is needed if the steroid injections do not work. It involves a small procedure under local anaesthetic. A slit is made in the mouth of the sheath to prevent the tendon catching at this point. Technically, this is achieved by the surgeon cutting the A1 pulley.

The operation is usually performed under local anaesthetic. The hand will be dressed with a supportive dressing that permits finger movement and light hand use. You will be discharged with specific instructions (“Hand Surgery”) relating to hand exercises and wound care. Your stitches will be removed two weeks after your operation.



Infection Any operation can be followed by infection and this would be treated with antibiotics.

Scar You will have a scar on the palm. This will be somewhat firm to touch and tender for 6-8 weeks. This can be helped by massaging the area firmly with the moisturizing cream.

Stiffness About 5% (1 in 20) of people are sensitive to hand surgery and their hand may become swollen, painful and stiff after any operation (algodystrophy). This problem cannot be predicted but will be watched for afterwards and treated with physiotherapy.

Nerve The nerves running to the fingers can be damaged during the surgery and cause numbness in part of the finger. This complication is very rare and the nerve would be repaired immediately.